



Basic Psychological Needs, Suicidal Ideation, and Risk for Suicidal Behavior in Young Adults

PETER C. BRITTON, PhD, KIMBERLY A. VAN ORDEN, PhD, JAMESON K. HIRSCH, PhD,
AND GEOFFREY C. WILLIAMS, MD, PhD

Associations between the satisfaction of basic psychological needs of autonomy, competence, and relatedness with current suicidal ideation and risk for suicidal behavior were examined. Two logistic regressions were conducted with a cross-sectional database of 440 university students to examine the association of need satisfaction with suicidal ideation and risk for suicidal behavior, while controlling for demographics and depressive symptoms. Suicidal ideation was reported by 15% of participants and 18% were found to be at risk for suicidal behavior. A one standard deviation increase in need satisfaction reduced the odds of suicidal ideation by 53%, OR (95% CI) = 0.47 (0.33–0.67), and the odds of being at risk for suicidal behavior by 50%, OR (95% CI) = 0.50 (0.37–0.69). Young adults whose basic psychological needs are met may be less likely to consider suicide and engage in suicidal behavior. Prospective research is needed to confirm these associations.

In 2009, suicide was the third leading cause of death among individuals ages 15 to 24

PETER C. BRITTON, Department of Veterans Affairs, Canandaigua Medical Center, VISN 2 Center of Excellence for Suicide Prevention, Canandaigua, NY, USA, and Center for the Study and Prevention of Suicide, Department of Psychiatry, University of Rochester Medical Center, Rochester, NY, USA; KIMBERLY A. VAN ORDEN, Center for the Study and Prevention of Suicide, Department of Psychiatry, University of Rochester Medical Center, Rochester, NY, USA; JAMESON K. HIRSCH, Department of Psychology, East Tennessee State University, Johnson City, TN, USA; GEOFFREY C. WILLIAMS, University of Rochester Medical Center, Rochester, NY, USA.

This paper was funded in part by a Career Development Award (K2CX000641) from the Department of Veterans Affairs Office of Research Development, Clinical Science Research and Development (CSR&D).

Address correspondence to Peter C. Britton, Center of Excellence for Suicide Prevention, Canandaigua VA Medical Center, 400 Fort Hill Ave., Canandaigua, NY 14424; e-mail: peter.britton@va.gov

and the second leading cause among those ages 25 to 34 (Centers for Disease Control & Prevention, 2012). The years of potential life lost in young adults represents significant losses to individuals, family, and friends, as well as to society. To inform prevention efforts in this population, researchers have started to examine protective factors that may reduce risk of suicidal ideation, suicide attempts, and suicide (Hirsch & Barton, 2011; Hirsch, Conner, & Duberstein, 2007; Hirsch, Wolford, Lalonde, Brunk, & Parker-Morris, 2009; Lamis, Ellis, Chumney, & Dula, 2009). Few studies have examined protective factors from the purview of a theory of human motivation, such as Self-determination Theory (SDT; Deci & Ryan, 1980; Ryan & Deci, 2002). Such theories suggest that individuals whose basic psychological needs are met may be less likely to consider and be at lower risk for suicidal behavior (e.g., suicide attempts and suicide; Britton, Patrick,

& Williams, 2011; Britton, Williams, & Conner, 2008).

SDT is an organismic theory that addresses the interaction between individual and environmental influences on human functioning and development. According to SDT, people are growth-oriented organisms that are intrinsically motivated to pursue and overcome challenges, and actualize their potential. However, SDT acknowledges that social-environmental influences can support or thwart people's natural tendency toward growth and development. SDT posits that three universal and basic psychological needs—for autonomy, competence, and relatedness—must be met in order for individuals to successfully pursue their innate interests and overcome the challenges they face (Deci & Ryan, 2000). People need to perceive themselves as *autonomous* and to experience their behavior as self-directed and congruent with their values and beliefs. They also need to have a sense of *competence* and believe that they can achieve their goals and excel in the activities they engage in. In addition, people are inherently social and require a sense of *relatedness* or acceptance and caring from others to remain motivated and engaged.

A subtheory of SDT, Basic Psychological Needs (BPN) theory, proposes that the universal and basic psychological needs for autonomy, competence, and relatedness are also directly associated with individuals' sense of well-being (Deci & Ryan, 2000). Research suggests that autonomy and competence are positively associated with indicators of physical and psychological well-being. In a meta-analysis using 184 data sets, higher levels of perceived autonomy and competence predicted higher levels of healthy behaviors (e.g. healthy eating, exercising), greater vitality, improved quality of life, greater satisfaction with life, fewer unhealthy behaviors (e.g., smoking, alcohol abuse), and lower levels of depression and anxiety (Ng et al., 2012). People who perceive themselves as autonomous and competent therefore have better physical and

mental health, which seems antithetical to the suicidal state. When their psychological needs are undermined, people have poorer physical and mental health, increasing their risk for negative outcomes such as suicide ideation and behavior. Thus, people who perceive that their basic psychological needs are supported are expected to be less likely to think about and engage in suicidal behavior.

The idea that psychological needs play a critical role in the development of suicidal thoughts and behavior provides the foundation for many theories of suicidal behavior (Joiner, 2005; Shneidman, 1996; Van Orden et al., 2010). Research shows that constructs related to these three basic psychological needs are associated with suicide-related outcomes, further supporting the importance of examining their relation with suicidal thoughts and behavior. Constructs related to autonomy and competence, such as locus of control, are associated with suicidal ideation and attempts in multiple populations including young adults (Beautrais, Joyce, & Mulder, 1999; Boor, 1979; Froyd & Perry, 1985). Burdensomeness, which is presumably associated with a low sense of competence (Joiner, 2005; Van Orden, Lynam, Hollar, & Joiner, 2006; Van Orden, Witte, Gordon, Bender, & Joiner, 2008), and deficits in problem solving have been found to be associated with suicidal ideation and attempts in a variety of populations including young adults (Dixon, Heppner, & Rudd, 1994; Schotte & Clum, 1982, 1987). There are also numerous findings supporting the association of low relatedness and higher rates of suicidal ideation and attempts (de Catanzaro, 1995; Groholt, Ekeberg, Wichstrom, & Haldorsen, 2000), and suicide in a variety of populations (Boardman, Grimbaldeston, Handley, Jones, & Willmott, 1999; Johansson, Sundquist, Johansson, & Bergman, 1997). A recent study of college students found that a higher order construct of "need satisfaction" (i.e., a composite score for all three needs

considered together) and each basic need separately moderated the relationship between negative life events and suicidal ideation (Rowe, Walker, Britton, & Hirsch, 2013).

The importance of psychological needs is also consistent with current thinking about the pathways through which social support could improve physical and mental health, thereby reducing risk for suicidal thoughts and behavior. Berkman, Glass, Brissette, and Seeman (2000) proposed one pathway whereby psychosocial factors increased positive health behaviors such as adherence to medical treatments, help-seeking behavior, maintaining a healthy diet, and exercising, resulting in improved health. This would suggest that individuals in environments that support autonomy, competence, or relatedness may be more likely to engage in healthy behavior that reduces their depression, hopelessness, and other factors that may increase their risk for suicidal thoughts and behavior. However, they also posited a direct psychological pathway whereby constructs including self-efficacy, coping effectiveness, and well-being directly influence health outcomes. Although not specifically stated by Berkman and colleagues, the hypothesis regarding a psychological pathway is consistent with the concept of psychological needs that must be satisfied to foster health. A direct influence on health presumes that people have needs for autonomy (i.e., self-efficacy), competence (i.e., coping effectiveness), and relatedness, just as they have needs for food, water, and shelter.

Building on previous research, in this study we examined the associations of basic need satisfaction with current suicidal ideation and risk for suicidal behavior in young adults. Given previous findings, we hypothesized that the satisfaction of basic needs would lower the odds of suicidal ideation and risk for suicidal behavior. Because individual needs are highly correlated but may have differential associations with each outcome (Johnston & Finney, 2010), explor-

atory analyses were also conducted for each individual need.

METHODS

Participants and Procedure

Participants in this study were 440 undergraduates recruited through general psychology courses who received course credit for their participation. Of the 440, 312 (71%) were female and 402 (91%) were White, non-Hispanic, with a mean age (*SD*) of 21.01 (6.10). After signing a statement of informed consent detailing the purpose, procedures, and goals of this study, participants completed a packet of self-report measures. All responses to suicide-related questions were screened for imminent suicide risk by study investigators to determine if immediate intervention was needed (Joiner, Walker, Rudd, & Jobes, 1999). After study completion, all participants were debriefed and provided telephone numbers for local mental health services. An institutional review board approved the project.

Measures

Suicidal Ideation. Current suicidal ideation was measured with item 9 from the Beck Depression Inventory-II (BDI-II; Beck, Steer, & Brown, 1996). The item was dichotomized to measure suicidal ideation (1 = "I have thoughts of killing myself but I would not carry them out" or "I would like to kill myself" or "I would kill myself if I had the chance") and no suicidal ideation (0 = "I don't have any thoughts of killing myself"). Validity for the BDI-II suicidal ideation item is well established (Beck, Brown, & Steer, 1997; Beck & Steer, 1991).

Risk for Suicidal Behavior. Risk for suicidal behavior was assessed with the Suicidal Behaviors Questionnaire Revised which has been validated with young adults (SBQ-R; Osman et al., 2001). The SBQ-R is comprised of four items ("Have you ever

thought about or attempted suicide?” “How often have you thought about killing yourself in the past year?” “Have you ever told someone that you were going to commit suicide, or that you might do it?” “How likely is it that you are going to commit suicide some day?”). Each item is scored with item-specific responses, ranging from 1–3 to 0–6. As recommended for nonclinical samples, a cutoff of seven or higher was used to indicate risk for suicidal behavior. Internal consistency for the SBQ-R was adequate ($\alpha = 0.78$, one-sided 95% CI ≥ 0.75).

Basic Psychological Needs. Psychological needs were assessed with the Basic Psychological Needs Scale (BPNS, Johnston & Finney, 2010), a 16-item self-report questionnaire that consists of three subscales assessing autonomy (three items; e.g., “I feel like I can pretty much be myself in my daily situations”), competence (six items, three reverse scored; e.g., “Most days I feel a sense of accomplishment from what I do”), and relatedness (seven items, two reverse scored; e.g., “I really like the people I interact with”), that has been validated with young adults. The BPNS is scored on a 7-point Likert scale with higher scores reflecting greater need support or fulfillment. Scores on the three scales were averaged and summed and a grand mean was calculated (Gagne, 2003). Internal consistency was good for the full scale ($\alpha = 0.89$, one-sided 95% CI ≥ 0.88), good for the

relatedness ($\alpha = 0.84$, one-sided 95% CI ≥ 0.82) subscale, and acceptable for the autonomy ($\alpha = 0.74$, one-sided 95% CI ≥ 0.71) and competence ($\alpha = 0.71$, one-sided 95% CI ≥ 0.67) subscales.

Depressive Symptoms. Depressive symptoms were assessed with the Beck Depression Inventory-II (BDI-II; (Beck, Steer, & Brown, 1996), a 21-item measure of cognitive and affective components of depression over the past 2 weeks. The BDI-II uses a 4-point Likert scale with higher scores indicative of increasingly severe symptoms. The BDI-II has well-established validity and reliability (Beck, Steer, Ball, & Ranieri, 1996; Beck, Steer, & Brown, 1996; Steer, Ball, Ranieri, & Beck, 1997), and has been shown to be valid in young adults (Robinson & Kelley, 1996). Prior to scoring, the suicidal ideation item was removed to prevent confounding with the outcomes. Internal consistency was excellent ($\alpha = 0.91$, one-sided 95% CI ≥ 0.90).

Statistical Analyses

Univariate analyses were used to examine independent associations with suicidal ideation and risk for suicidal behavior. Two multivariate logistic regressions were conducted; the first predicted current suicidal ideation, and the second predicted risk for suicidal behavior. All variables were entered in the multivariate regression models using forced entry. Variables included

TABLE 1
Basic Psychological Needs and Suicidal Ideation in Young Adults (N = 440)

	NonIdeators N = 375 n (%)	Ideators N = 65 n (%)	Univariate Analyses OR (95% CI)	Multivariate Analysis OR (95% CI)
Female	273 (73%) M (SD)	39 (60%) M (SD)	0.56 (0.32–0.97)	0.61 (0.32–1.18)
Age	21.10 (6.38)	20.49 (4.16)	0.98 (0.93–1.03)	0.97 (0.92–1.02)
Depression ^a	1.97 (2.05)	5.25 (3.41)	3.10 (2.33–4.12)	2.25 (1.63–3.11)
BPNS ^a	5.54 (0.81)	4.50 (0.88)	0.31 (0.23–0.43)	0.47 (0.33–0.67)

BPNS, Basic Psychological Needs Scale. ^aConverted to z scores in univariate and multivariate analyses.

were standard demographic variables (i.e., gender and age), depressive symptoms to ensure that they did not explain the results (Eisenberg, Gollust, Golberstein, & Hefner, 2007; Konick & Gutierrez, 2005; Van Orden et al., 2008), and the basic needs. Basic needs and depression scores were converted to *z*-scores to improve interpretation. The Hosmer–Lemeshow goodness-of-fit statistic was used to evaluate model fit (Hosmer & Lemeshow, 1989), and odds ratios with 95% confidence intervals were derived using the method of maximum likelihood. Exploratory univariate and multivariate logistic regressions were conducted for each individual need, using the same procedures described earlier.

RESULTS

Suicidal Ideation

Of the 440 participants, 65 (15%) reported current suicidal ideation. In univariate analyses, basic need satisfaction was associated with suicidal ideation (Table 1). The multivariate model for suicidal ideation provided adequate fit, $\chi^2(8) = 2.56, p = 0.96$. A one standard deviation increase in need satisfaction decreased the odds of suicidal ideation by 53%, OR (95% CI) = 0.47 (0.33–0.67), and a one standard deviation increase in depressive symptoms more than doubled the odds of suicidal ideation, OR (95% CI) = 2.25 (1.63–3.11).

In exploratory univariate analyses, autonomy, competence, and relatedness were associated with suicidal ideation (Table 2). The multivariate model examining autonomy and suicidal ideation provided adequate fit, $\chi^2(8) = 5.74, p = 0.68$. A one standard deviation increase in perceived autonomy reduced the odds of suicidal ideation by 45%, OR (95% CI) = 0.55 (0.40–0.76), and a one standard deviation increase in depressive symptoms more than doubled the odds of suicidal ideation, OR (95% CI) = 2.54 (1.85–3.47). The multivariate model examining the association between

TABLE 2
Individual Psychological Needs and Suicidal Ideation in Young Adults (N = 440)

	Nonideators N = 375		Ideators N = 65		Univariate Analyses OR (95% CI)	Autonomy Multivariate Analysis OR (95% CI)	Competence Multivariate Analysis OR (95% CI)	Relatedness Multivariate Analysis OR (95% CI)
	M (SD)	M (SD)	M (SD)	M (SD)				
Female	—	—	—	—	—	0.61 (0.32–1.16)	0.56 (0.29–1.07)	0.60 (0.31–1.13)
Age	—	—	—	—	—	0.97 (0.92–1.03)	0.97 (0.92–1.02)	0.95 (0.90–1.01)
Depression ^a	—	—	—	—	—	2.54 (1.85–3.47)	2.24 (1.60–3.11)	2.65 (1.94–3.60)
Autonomy ^a	5.60 (1.02)	—	4.47 (1.21)	—	0.39 (0.29–0.51)	0.55 (0.40–0.76)	—	—
Competence ^a	5.24 (0.92)	—	4.14 (0.93)	—	0.30 (0.21–0.41)	—	—	—
Relatedness ^a	5.78 (0.90)	—	4.89 (1.05)	—	0.44 (0.34–0.57)	0.46 (0.31–0.68)	—	0.62 (0.45–0.84)

^aConverted to *z* scores in univariate and multivariate analyses.

TABLE 3
Individual Psychological Needs and Risk for Suicidal Behavior in Young Adults (N = 440)

	Low Risk N = 360 n (%)	High Risk N = 80 n (%)	Univariate Analyses OR (95% CI)	Multivariate Analysis OR (95% CI)
Female	258 (72%)	54 (68%)	0.82 (0.49–1.38)	0.98 (0.55–1.76)
	<i>M (SD)</i>	<i>M (SD)</i>		
Age	20.98 (6.41)	21.16 (4.45)	1.00 (0.97–1.04)	1.00 (0.96–1.05)
Depression ^a	2.04 (2.17)	4.35 (3.32)	2.22 (1.74–2.82)	1.59 (1.20–2.10)
BPNS ^a	5.54 (0.79)	4.69 (1.02)	0.39 (0.30–0.52)	0.50 (0.39–0.69)

BPNS, Basic Psychological Needs Scale. ^aConverted to *z* scores in univariate and multivariate analyses.

competence and suicidal ideation provided adequate fit, $\chi^2(8) = 6.69$, $p = 0.57$. A one standard deviation increase in perceived competence decreased the odds of suicidal ideation by 54%, OR (95% CI) = 0.46 (0.31–0.68), and a one standard deviation increase in depressive symptoms more than doubled the odds of suicidal ideation, OR (95% CI) = 2.24 (1.60–3.11). The multivariate model examining relatedness and suicidal ideation also provided adequate fit, $\chi^2(8) = 4.23$, $p = 0.84$. A one standard deviation increase in perceived relatedness reduced the odds suicidal ideation by 38%, OR (95% CI) = 0.62 (0.45–0.84), and a one standard deviation increase in depressive symptoms more than doubled the odds of suicidal ideation, OR (95% CI) = 2.65 (1.94–3.60).

Risk for Suicidal Behavior

Of the 440 participants, 80 (18%) scored seven or above on the SBQ-R indicating risk for suicidal behavior. In univariate analyses, basic need satisfaction was associated with increased risk for suicidal behavior (see Table 3). The multivariate model for risk for suicidal behavior provided adequate fit, $\chi^2(8) = 13.66$, $p = 0.09$. A one standard deviation increase in need satisfaction reduced the odds of being at risk for suicidal behavior by 50%, OR (95% CI) = 0.50 (0.37–0.69), and a one standard deviation increase in depressive symptoms increased the odds of being at risk for

suicidal behavior by 59%, OR (95% CI) = 1.59 (1.20–2.10).

In exploratory univariate and multivariate analyses, autonomy, competence, and relatedness were associated with risk for suicidal behavior (Table 2). The multivariate model examining autonomy and risk for suicidal behavior provided adequate fit, $\chi^2(8) = 6.30$, $p = 0.61$. A one standard deviation increase in perceived autonomy reduced the odds of being at risk for suicidal behavior by 33%, OR (95% CI) = 0.67 (0.51–0.88), and a one standard deviation increase in depressive symptoms increased the odds of being at risk for suicidal behavior by 87%, OR (95% CI) = 1.87 (1.44–2.44). The multivariate model examining competence and risk for suicidal behavior provided adequate fit, $\chi^2(8) = 7.03$, $p = 0.53$. A one standard deviation increase in perceived competence reduced the odds of being at risk for suicidal behavior by 51%, OR (95% CI) = 0.49 (0.35–0.68), and a one standard deviation increase in depressive symptoms increased the odds of being at risk for suicidal behavior by 56%, OR (95% CI) = 1.56 (1.17–2.08). The multivariate model examining relatedness and suicide attempts also provided adequate fit, $\chi^2(8) = 5.93$, $p = 0.66$. A one standard deviation increase in perceived relatedness reduced the odds of being at risk for suicidal behavior by 42%, OR (95% CI) = 0.58 (0.44–0.76), and a one standard deviation increase in depressive symptoms

TABLE 4
Individual Psychological Needs and Risk for Suicidal Behavior in Young Adults (N = 440)

	Low Risk		High Risk		Univariate Analyses		Autonomy Multivariate		Competence Multivariate		Relatedness Multivariate	
	N = 360	N = 80	M (SD)	M (SD)	OR (95% CI)	Analysis OR (95% CI)	Analysis OR (95% CI)	Analysis OR (95% CI)	Analysis OR (95% CI)	Analysis OR (95% CI)	Analysis OR (95% CI)	
Female	—	—	—	—	—	0.95 (0.54–1.68)	—	0.91 (0.51–1.62)	—	0.98 (0.55–1.74)	—	
Age	—	—	—	—	—	1.00 (0.96–1.04)	—	1.00 (0.96–1.05)	—	0.99 (0.95–1.04)	—	
Depression ^a	—	—	—	—	—	1.87 (1.44–2.44)	—	1.56 (1.17–2.08)	—	1.79 (1.38–2.32)	—	
Autonomy ^a	5.58 (1.02)	4.78 (1.32)	—	—	0.51 (0.40–0.65)	0.67 (0.51–0.88)	—	—	—	—	—	
Competence ^a	5.24 (0.90)	4.33 (1.08)	—	—	0.38 (0.28–0.50)	—	—	—	—	—	—	
Relatedness ^a	5.79 (0.88)	4.98 (1.09)	—	—	0.46 (0.36–0.58)	—	—	0.49 (0.35–0.68)	—	0.58 (0.44–0.76)	—	

^aConverted to z scores in univariate and multivariate analyses.

increased the odds of being at risk for suicidal behavior by 79%, OR (95% CI) = 1.79 (1.38–2.32).

DISCUSSION

In this study we examined the association of the basic psychological needs of autonomy, competence, and relatedness with current suicidal ideation and risk for suicidal behavior in a young adult sample. Our primary hypothesis was supported as total need satisfaction was associated with lower odds of suicidal ideation and risk for suicidal behavior. Exploratory analyses provided further support, suggesting that the satisfaction of the needs for autonomy, competence, and relatedness were also independently associated with lower odds of suicidal ideation and risk for suicidal behavior.

These findings add to a growing literature using BPN theory to better understand the factors that reduce risk for suicidal ideation and suicidal behavior in young adults. BPN theory posits that the satisfaction of the basic psychological needs for autonomy, competence, and relatedness are associated with an increased sense of well-being that presumably lowers risk for suicidal thoughts and behavior. Previous findings show that support of these basic needs serve as a buffer between negative life events and suicidal ideation (Rowe et al., 2013). These findings support that satisfaction of these basic needs provides protection against suicidal ideation, and suggest it may also protect against suicidal behavior. This advancement is critical as previous investigations have focused solely on suicidal thoughts.

If they are replicated in prospective studies, these findings may have important implications for the prevention of suicidal thoughts and behavior in young adults. Interventions based on SDT principles are associated with increased engagement in healthy behaviors such as smoking cessation and diabetes management (Williams, McGregor, King, Nelson, & Glasgow,

2005; Williams, McGregor, Sharp, Kouides, et al., 2006; Williams, McGregor, Sharp, Levesque, et al., 2006; Williams, McGregor, Zeldman, Freedman, & Deci, 2004; Williams, Niemiec, Patrick, Ryan, & Deci, 2009). Prospective studies supporting the associations between basic psychological needs and suicidal ideation and suicidal behavior would support the application of interventions that are congruent with BPN theory such as motivational interviewing to suicide prevention (Britton, Conner, & Maisto, 2012; Britton et al., 2008, 2011).

A number of limitations must be considered when interpreting these findings. Data were cross-sectional, preventing causality and directionality from being established. For example, need satisfaction may reduce risk for suicidal ideation, but suicidal thoughts could also negatively impact an individual's sense of autonomy, competence, and relatedness. Potentially important contributors to suicidal thoughts and behavior in young adults, such as alcohol and drug use, were not available for analysis and

should be included in future studies. Data were collected from a sample of predominantly white, female undergraduate students, reducing generalizability.

Nonetheless, this study suggests that satisfaction of basic psychological needs may lower the odds that young adults consider or engage in suicidal behavior. These findings also support the notion that SDT-based constructs such as supporting or thwarting the satisfaction of basic psychological needs, autonomy, intrinsic motivation, and goal pursuit and attainment may have important implications for understanding and preventing suicidal thoughts and behavior in this population (Bureau, Mageau, Vallerand, Rousseau, & Otis, 2012; Deci & Ryan, 2000). Future research utilizing prospective designs should examine the associations among the basic psychological needs, possible mechanisms (i.e., help-seeking, treatment engagement, intrinsic motivation, vitality), and suicidal ideation and behavior in young adults.

REFERENCES

- BEAUTRAIS, A. L., JOYCE, P. R., & MULDER, R. T. (1999). Personality traits and cognitive styles as risk factors for serious suicide attempts among young people. *Suicide and Life-Threatening Behavior, 29*, 37–47.
- BECK, A. T., BROWN, G. K., & STEER, R. A. (1997). Psychometric characteristics of the Scale for Suicide Ideation with psychiatric outpatients. *Behaviour Research and Therapy, 35*, 1039–1046.
- BECK, A. T., & STEER, R. A. (1991). *Manual for the Beck Scale for Suicide Ideation*. San Antonio, TX: Psychological Corporation.
- BECK, A. T., STEER, R. A., BALL, R., & RANIERI, W. F. (1996). Comparison of Beck Depression Inventories-IA and -II in psychiatric outpatients. *Journal of Personality Assessment, 67*, 588–597.
- BECK, A. T., STEER, R. A., & BROWN, G. K. (1996). *Manual for the Beck Depression Inventory-II*. San Antonio, TX: Psychological Corporation.
- BERKMAN, L. F., GLASS, T., BRISSETTE, I., & SEEMAN, T. E. (2000). From social integration to health: Durkheim in the new millennium. *Social Science & Medicine, 51*, 843–857.
- BOARDMAN, A. P., GRIMBALDESTON, A. H., HANDLEY, C., JONES, P. W., & WILLMOTT, S. (1999). The North Staffordshire suicide study: A case-control study of suicide in one health district. *Psychological Medicine, 29*, 27–33.
- BOOR, M. (1979). Relationship of internal-external control and United States suicide rates, 1973-1976. *Journal of Clinical Psychology, 35*, 513–516.
- BRITTON, P. C., CONNER, K. R., & MAISTO, S. A. (2012). An open trial of motivational interviewing to address suicidal ideation with hospitalized veterans. *Journal of Clinical Psychology, 68*(9), 961–971.
- BRITTON, P. C., PATRICK, H., & WILLIAMS, G. C. (2011). Motivational interviewing, self-determination theory, and cognitive behavioral therapy to prevent suicidal behavior. *Journal of Cognitive Behavioral Practice, 18*, 16–27.
- BRITTON, P. C., WILLIAMS, G. C., & CONNER, K. R. (2008). Self-determination theory, motivational interviewing, and the treatment of clients with acute suicidal ideation. *Journal of Clinical Psychology, 64*, 52–66.

- BUREAU, J. S., MAGEAU, G. A., VALLERAND, R. J., ROUSSEAU, F. L., & OTIS, J. (2012). Self-determination: A buffer against suicide ideation. *Suicide and Life-Threatening Behavior, 42*, 377–393.
- DE CATANZARO, D. (1995). Reproductive status, family interactions, and suicidal ideation: Surveys of the general public and high-risk groups. *Ethology & Sociobiology, 16*, 385–394.
- Centers for Disease Control and Prevention. (2012). *Wisqars injury mortality reports 1999–2009*. Retrieved April 12, 2012, from <http://www.cdc.gov.ezpmminer.urmc.rochester.edu/ncipc/wisqars/>.
- DECI, E. L., & RYAN, R. M. (1980). Self-determination theory: When mind mediates behavior. *Journal of Mind and Behavior, 1*, 33–43.
- DECI, E. L., & RYAN, R. M. (2000). The “what” and “why” of goal pursuits: Human needs and the self-determination of behavior. *Psychological Inquiry, 11*, 227–268.
- DIXON, W. A., HEPPNER, P. P., & RUDD, M. D. (1994). Problem-solving appraisal, hopelessness, and suicide ideation: Evidence for a mediational model. *Journal of Counseling Psychology, 41*, 91–98.
- EISENBERG, D., GOLLUST, S. E., GOLBERSTEIN, E., & HEFNER, J. L. (2007). Prevalence and correlates of depression, anxiety, and suicidality among university students. *American Journal of Orthopsychiatry, 77*, 534–542.
- FROYD, J., & PERRY, N. (1985). Relationships among locus of control, coronary-prone behavior, and suicidal ideation. *Psychological Reports, 57*, 1155–1158.
- GAGNE, M. (2003). The role of autonomy support and autonomy orientation in prosocial behavior engagement. *Motivation and Emotion, 27*, 199–223.
- GROHOLT, B., EKEBERG, O., WICHSTROM, L., & HALDORSEN, T. (2000). Young suicide attempters: A comparison between a clinical and an epidemiological sample. *Journal of the American Academy of Child and Adolescent Psychiatry, 39*, 868–875.
- HIRSCH, J. K., & BARTON, A. L. (2011). Positive social support, negative social exchanges, and suicidal behavior in college students. *Journal of American College Health, 59*, 393–398.
- HIRSCH, J. K., CONNER, K. R., & DUBERSTEIN, P. R. (2007). Optimism and suicide ideation among young adult college students. *Archives of Suicide Research, 11*, 177–185.
- HIRSCH, J. K., WOLFORD, K., LALONDE, S. M., BRUNK, L., & PARKER-MORRIS, A. (2009). Optimistic explanatory style as a moderator of the association between negative life events and suicide ideation. *Crisis, 30*, 48–53.
- HOSMER, D. W., & LEMESHOW, S. (1989). *Applied logistic regression*. New York: Wiley.
- JOHANSSON, L. M., SUNDQUIST, J., JOHANSSON, S. E., & BERGMAN, B. (1997). Ethnicity, social factors, illness and suicide: A follow-up study of a random sample of the Swedish population. *Acta Psychiatrica Scandinavica, 95*, 125–131.
- JOHNSTON, M. M., & FINNEY, S. J. (2010). Measuring basic needs satisfaction: Evaluating previous research and conducting new psychometric evaluations of the basic needs satisfaction in general scale. *Contemporary Educational Psychology, 35*, 280–296.
- JOINER, T. E., JR. (2005). *Why people die by suicide*. Cambridge: Harvard University Press.
- JOINER, T. E., JR., WALKER, R. L., RUDD, M. D., & JOBES, D. A. (1999). Scientizing and routinizing the assessment of suicidality in outpatient practice. *Professional Psychology, 30*, 447–453.
- KONICK, L. C., & GUTIERREZ, P. M. (2005). Testing a model of suicide ideation in college students. *Suicide and Life-Threatening Behavior, 35*, 181–192.
- LAMIS, D. A., ELLIS, J. B., CHUMNEY, F. L., & DULA, C. S. (2009). Reasons for living and alcohol use among college students. *Death Studies, 33*, 277–286.
- NG, J. Y. Y., NTOUMANIS, N., THOGENSEN-NTOUMANI, C., DECI, E. L., RYAN, R. M., DUDA, J. L., ET AL. (2012). Self-determination theory applied to health contexts: A meta-analysis. *Perspectives on Psychological Science, 7*, 325–340.
- OSMAN, A., BAGGE, C. L., GUTIERREZ, P. M., KONICK, L. C., KOPPER, B. A., & BARRIOS, F. X. (2001). The Suicidal Behaviors Questionnaire-Revised (SBQ-R): Validation with clinical and nonclinical samples. *Assessment, 8*, 443–454.
- ROBINSON, B. E., & KELLEY, L. (1996). Concurrent validity of the Beck Depression Inventory as a measure of depression. *Psychological Reports, 79*, 929–930.
- ROWE, B. S., WALKER, K. L., BRITTON, P. C., & HIRSCH, J. K. (2013). The relationship between negative life events and suicidal behavior: Moderating role of basic psychological needs. *Crisis, 34*, 233–241.
- RYAN, R. M., & DECI, E. L. (2002). Overview of self-determination theory: An organismic-dialectical perspective. In E. L. Deci & R. M. Ryan (Eds.), *Handbook of self-determination research* (pp. 3–33). Rochester, NY: University of Rochester Press.
- SCHOTTE, D. E., & CLUM, G. A. (1982). Suicide ideation in a college population: A test of a model. *Journal of Consulting and Clinical Psychology, 50*, 690–696.

- SCHOTTE, D. E., & CLUM, G. A. (1987). Problem-solving skills in suicidal psychiatric patients. *Journal of Consulting and Clinical Psychology, 55*, 49–54.
- SHNEIDMAN, E. S. (1996). *The suicidal mind*. New York: Oxford University Press.
- STEER, R. A., BALL, R., RANIERI, W. F., & BECK, A. T. (1997). Further evidence for the construct validity of the beck Depression Inventory-II with psychiatric outpatients. *Psychological Reports, 80*, 443–446.
- VAN ORDEN, K. A., LYNAM, M. E., HOLLAR, D., & JOINER, T. E., JR. (2006). Perceived burdenomeness as an indicator of suicidal symptoms. *Cognitive Therapy and Research, 30*, 457–467.
- VAN ORDEN, K. A., WITTE, T. K., CUKROWICZ, K. C., BRAITHWAITE, S. R., SELBY, E. A., & JOINER, T. E., JR. (2010). The interpersonal theory of suicide. *Psychological Review, 117*, 575–600.
- VAN ORDEN, K. A., WITTE, T. K., GORDON, K. H., BENDER, T. W., & JOINER, T. E., JR. (2008). Suicidal desire and the capability for suicide: Tests of the interpersonal-psychological theory of suicidal behavior among adults. *Journal of Consulting and Clinical Psychology, 76*, 72–83.
- WILLIAMS, G. C., MCGREGOR, H. A., KING, D., NELSON, C. C., & GLASGOW, R. E. (2005). Variation in perceived competence, glycemic control, and patient satisfaction: Relationship to autonomy support from physicians. *Patient Education & Counseling, 57*, 39–45.
- WILLIAMS, G. C., MCGREGOR, H. A., SHARP, D., KOUIDES, R. W., LEVESQUE, C., RYAN, R. M., ET AL. (2006). A self-determination multiple risk intervention trial to improve smokers' health. *Journal of General Internal Medicine, 21*, 1288–1294.
- WILLIAMS, G. C., MCGREGOR, H. A., SHARP, D., LEVESQUE, C., KOUIDES, R. W., RYAN, R. M., ET AL. (2006). Testing a self-determination theory intervention for motivating tobacco cessation: Supporting autonomy and competence in a clinical trial. *Health Psychology, 25*, 91–101.
- WILLIAMS, G. C., MCGREGOR, H. A., ZELDMAN, A., FREEDMAN, Z. R., & DECI, E. L. (2004). Testing a self-determination theory process model for promoting glycemic control through diabetes self-management. *Health Psychology, 23*, 58–66.
- WILLIAMS, G. C., NIEMIEC, C. P., PATRICK, H., RYAN, R. M., & DECI, E. L. (2009). The importance of supporting autonomy and perceived competence in facilitating long-term tobacco abstinence. *Annals of Behavioral Medicine, 37*, 315–324.

Manuscript Received: November 18, 2012
 Revision Accepted: September 17, 2013