


This article was downloaded by: [Curtin University Library]

On: 20 May 2015, At: 18:35

Publisher: Routledge

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



[Click for updates](#)

Psychology, Health & Medicine

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/cphm20>

Motivational processes and well-being in cardiac rehabilitation: a self-determination theory perspective

Rachel Jane Rahman^a, Joanne Hudson^b, Cecilie Thøgersen-Ntoumani^c & Jonathan H. Doust^d

^a Department of Psychology, Aberystwyth University, Penbryn 5, Penglais Campus, Aberystwyth SY23 3UX, Ceredigion, UK

^b Department of Sport and Exercise Science, Aberystwyth University, Carwyn James Building, Penglais Campus, Aberystwyth SY23 3FD, Ceredigion, UK

^c School of Sport and Exercise Sciences, College of Life and Environmental Sciences, University of Birmingham, Edgbaston B15 2TT, UK

^d Chelsea School, Brighton University, Hillbrow, Gaudick Road, Eastbourne BN20 7SR, UK

Published online: 10 Mar 2015.

To cite this article: Rachel Jane Rahman, Joanne Hudson, Cecilie Thøgersen-Ntoumani & Jonathan H. Doust (2015) Motivational processes and well-being in cardiac rehabilitation: a self-determination theory perspective, *Psychology, Health & Medicine*, 20:5, 518-529, DOI: [10.1080/13548506.2015.1017509](https://doi.org/10.1080/13548506.2015.1017509)


To link to this article: <http://dx.doi.org/10.1080/13548506.2015.1017509>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

Motivational processes and well-being in cardiac rehabilitation: a self-determination theory perspective

Rachel Jane Rahman^{a*} , Joanne Hudson^{b,1}, Cecilie Thøgersen-Ntoumani^{c,2} and Jonathan H. Doust^d

^aDepartment of Psychology, Aberystwyth University, Penbryn 5, Penglais Campus, Aberystwyth SY23 3UX, Ceredigion, UK; ^bDepartment of Sport and Exercise Science, Aberystwyth University, Carwyn James Building, Penglais Campus, Aberystwyth SY23 3FD, Ceredigion, UK; ^cSchool of Sport and Exercise Sciences, College of Life and Environmental Sciences, University of Birmingham, Edgbaston B15 2TT, UK; ^dChelsea School, Brighton University, Hillbrow, Gaudick Road, Eastbourne BN20 7SR, UK

(Received 11 August 2014; accepted 3 February 2015)

This research examined the processes underpinning changes in psychological well-being and behavioural regulation in cardiac rehabilitation (CR) patients using self-determination theory (SDT). A repeated measures design was used to identify the longitudinal relationships between SDT variables, psychological well-being and exercise behaviour during and following a structured CR programme. Participants were 389 cardiac patients (aged 36–84 years; $M_{\text{age}} = 64 \pm 9$ years; 34.3% female) referred to a 12-week-supervised CR programme. Psychological need satisfaction, behavioural regulation, health-related quality of life, physical self-worth, anxiety and depression were measured at programme entry, exit and six month post-programme. During the programme, increases in autonomy satisfaction predicted positive changes in behavioural regulation, and improvements in competence and relatedness satisfaction predicted improvements in behavioural regulation and well-being. Competence satisfaction also positively predicted habitual physical activity. Decreases in external regulation and increases in intrinsic motivation predicted improvements in physical self-worth and physical well-being, respectively. Significant longitudinal relationships were identified whereby changes during the programme predicted changes in habitual physical activity and the mental quality of life from exit to six month follow-up. Findings provide insight into the factors explaining psychological changes seen during CR. They highlight the importance of increasing patients' perceptions of psychological need satisfaction and self-determined motivation to improve well-being during the structured component of a CR programme and longer term physical activity.

Keywords: self-determination theory; cardiac rehabilitation; well-being; psychological need satisfaction; behavioural regulation

Cardiac rehabilitation (CR) programmes are an essential component of care for patients who have experienced MI, cardiovascular disease and pre-/post-cardiac surgery (National Institute for Clinical Excellence [NICE], 2013). Typically, programmes involve a

*Corresponding author. Email: rjr@aber.ac.uk

¹Current Affiliation: Carnegie Faculty, School of Sport, Leeds Beckett University, Headingley Campus, Leeds LS6 3QS, West Yorkshire, UK.

²Current Affiliation: Faculty of Health Sciences, School of Psychology & Speech Pathology, Curtin University, GPO Box U1987, Perth, Western Australia 6845, Australia.

multidisciplinary team offering exercise training, education and counselling, with the aim of improving physical functioning, symptoms and quality of life (NICE, 2013).

CR programmes have been found to reduce deaths in cardiac patients by 27% (British Heart Foundation [BHF], 2010); improve exercise capacity, lipid, lipoprotein and blood glucose levels; and reduce body weight, systolic and diastolic blood pressure (Balady, Fletcher, & Froelicher, 1994; Bjarnason-Wehrens et al., 2007). Research has also revealed psychological benefits in the form of improved anxiety, depression and quality of life (Yohannes, Doherty, Bundy, & Yalfani, 2010). This is important given that cardiac patients report elevated levels of anxiety and depression and poor quality of life, contributing to increased risks of secondary cardiac events (Frasure-Smith & Lespérance, 2005, 2008). Despite this, only 40% of MI patients participate in CR programmes, highlighting the need to understand exercise motivation in CR patients (BHF, 2010).

Studies have drawn on psychological theory to understand motivations underpinning exercise behaviour and participation in CR. One theory demonstrating relevant application is self-determination theory (SDT; Deci & Ryan, 1985). SDT is concerned with the processes involved in behavioural regulation and their associated cognitive, affective and behavioural outcomes. According to SDT, individuals' self-determined regulation lies along a continuum of five increasingly internalised regulations. Amotivation signifies a lack of motivation, external regulation refers to a drive as a result of pressure from an external source, introjected regulation refers to feeling moved to engage in a behaviour but not truly accepting its value, and identified regulation refers to feeling motivated through accepting the values of the behaviour in question. Finally, intrinsic motivation signifies the most self-determined form of regulation where individuals engage in behaviour for the inherent pleasure of doing so. According to SDT, self-determined regulation leads to increased persistence, well-being and likelihood to maintain behaviour (Deci & Ryan, 2002).

SDT further proposes that self-determined regulation and well-being result from the satisfaction of three innate psychological needs: Autonomy (the need to feel volitional in one's actions), competence (the need to feel able to affect outcomes) and relatedness (the need to have supportive relationships; Wilson, Rodgers, Blanchard, & Gessell, 2003).

SDT has demonstrated relevance in exercise and health care contexts. As such, self-determined regulations lead to better mental health and stronger physical activity intentions (Rouse, Ntoumanis, Duda, Jolly, & Williams, 2011) and improvements in psychological outcomes in exercise referral patients (Rahman, Thogersen-Ntoumani, Thatcher, & Doust, 2011). These associations have been confirmed by meta-analysis (Ng et al., 2012).

More specifically, in CR, Russell and Bray (2009) used SDT to predict exercise behaviour in a cross-sectional and prospective study of CR outpatients ($N = 68$, $M_{\text{age}} = 64.9$ years) with competence satisfaction demonstrating a key role as a predictor of self-determined regulation and exercise behaviour up to six weeks following CR. Extending these findings, Sweet, Tulloch, Fortier, Pipe, and Reid (2011) demonstrated that self-determined individuals were more likely to maintain exercise behaviour up to 24 months following CR ($N = 251$, 79% male, $M_{\text{age}} = 61.4$ years).

These studies illustrate the applicability of SDT for predicting short- and long-term exercise behaviour; however, they did not examine relationships between SDT variables and psychological outcomes in a CR context. The aim of this research was therefore to use SDT to explore whether changes in psychological need satisfaction predicted changes in behavioural regulation over the duration of the programme and whether changes in psychological need satisfaction and behavioural regulation during the

programme predicted changes in psychological well-being during the programme and from exit to six month follow-up.

It was hypothesised that increases in psychological need satisfaction would predict improvements in self-determined regulation and decreases in more controlled regulations during the programme (entry to exit) and that increases in psychological need satisfaction and self-determined regulation and decreases in controlled forms of regulation during the programme (entry to exit) would predict improvements in psychological well-being and habitual physical activity during the programme (entry to exit) and beyond (exit to six months follow-up).

Methods

Participants

577 participants were referred to the CR scheme over three years (age range 18–87 years; $M_{\text{age}} = 64 \pm 10$ years; 35.3% female). 484 were invited to start classes during the research period and 389 consented (age range 36–84 years; $M_{\text{age}} = 64 \pm 9$ years; 34.3% female). No significant differences in age ($t(721) = .78, p > .05$) or gender ($Z(1) = -.44, p > .05$) were identified between the total referral group and the research participants. Figure 1 presents a flow diagram of participation which illustrates that not all of the 389 consenting participants returned each questionnaire at each time point. 243 completed the programme during the research period. Many did not provide a reason for dropout (48%), four died during the programme (non-programme related), other reasons included ill-health, family or work commitments, moving and lack of time. Comparisons of programme completers and non-completers using a Mann–Whitney test demonstrated a significant difference in anxiety ($U(248) = 5392.5, p < .05$), depression ($U(248) = 5294.0, p < .05$) and mental component summary (MCS) of the SF-36v2 (see below; $U(245) = 5645.0, p < .05$) with completers reporting significantly higher levels of well-being.

Measures

Behavioural Regulation in Exercise Questionnaire-2 (BREQ-2; Markland & Tobin, 2004a) measured participants' exercise regulation and comprised 19 items [five-point scale ranging from 0 (*Not true for me*) to 4 (*Very true for me*)] with 5 subscales representing the motivational regulations. Reliability analyses show high Cronbach's alphas ranging from .73 to .86 for subscales (Markland & Tobin, 2004a).

Psychological Need Satisfaction Scale (PNSS; Markland & Tobin, 2004b) comprises nine items (three for each need) measuring psychological need satisfaction in an exercise class context. The response scale matches that of the BREQ-2. Cronbach's alphas range between .59 and .72 (Markland & Tobin, 2004b).

Hospital Anxiety and Depression Scale (HADS; Zigmond & Snaith, 1983) is a 14-item questionnaire [four-point scale; scores ranging from 0 to 21] consisting of subscales for anxiety and depression. Both subscales report alpha values exceeding .90 (Moorey et al., 1991).

Short Form-36version2 (SF-36v2; Wade, Snow, Kosinski, & Gandek, 1993, 2002) measured health-related quality of life. The 36 items measure physical and social functioning, role limitations due to physical and emotional problems, mental health, vitality, pain and general health. Response scales vary by question. Scores form two summaries: The physical component summary (PCS) and the MCS. Cronbach's alphas for the subscales range from .80 to .95 (Jenkinson, Stewart-Brown, Petersen, & Paice, 1999).




Figure 1. Flow diagram of participant involvement in research.

Baecke's Questionnaire of Habitual Physical Activity (BeackeHPA; Baecke, Burema, & Frijters, 1982) comprises 16 items that constitute three subscales: Activity at work, sport and leisure. Response format varies by question and responses are combined,

producing an overall habitual activity score. Cronbach's alphas for the total score are reported as .77 (Florindo, do Rosario Dias, & Latorre, 2003).

Physical Self-Perception Profile (PSPP; Fox & Corbin, 1989) comprises 5 subscales. The physical self-worth subscale used in this study assesses an individual's general feelings about their physical self. The question format employs a four-point forced-choice scale in which two alternative statements are provided and individuals select the most representative statement before indicating the degree to which the statement is true for them. Coefficient alphas for the subscale exceed .80 (Fox & Corbin, 1989).

The CR programme

A 12-week-supervised Phase III (structured exercise training and education) and IV (maintenance) CR programme was held in six council-owned leisure centres across mid-Wales. Adult participants with any heart condition were referred by health professionals or could self-refer. They included those who had suffered an MI, were awaiting/recovering from heart surgery and those with heart disease, heart failure, stable angina or controlled arrhythmias. Anyone with uncontrolled hypertension, unstable angina or uncontrolled arrhythmias was not eligible.

Patients were invited to a consultation where a cardiac nurse assessed their medical history, they were briefed about the research, and their participation was requested. Participants provided informed consent and were invited to attend an induction followed by twice weekly 60 min exercise classes: One gym based and one circuit session. Qualified exercise professionals taught participants how to exercise safely and gradually increase exercise intensity. Both exercise sessions included cardiovascular and strength-based exercise with warm up-/warm down protocols of approximately 10 min within the hour. Participants wore heart rate monitors and were encouraged to work between 60 and 75% of their maximum heart rate, as determined by the exercise professionals. On completion of the programme, participants were offered a six month maintenance package, which entitled them to access exercise facilities at reduced cost.

Procedures

Ethical approval for the study was obtained from an NHS ethics Committee. The SF-36v2, PSPP and the HADS were posted for completion prior to commencing classes. Participants met with a researcher during their first class and completed the BaeckeHPA. The BREQ-2, the PNSS, a covering letter and a prepaid return envelope were sent to participants following their first class. Following their final class, participants met with the researcher to complete the BaeckeHPA and were asked to complete all other measures independently. Those who completed the programme were also contacted six months later and asked to complete all measures again.

Data analysis

Assumptions for multicollinearity, independence of outcome variables, independent errors homoscedasticity, normally distributed errors and linearity were tested. Data were not normally distributed and remained so following log and root transformations. Thus, original raw data were retained. An intent-to-treat analysis was utilised where baseline data were carried forward to missing time points to provide a conservative effect of

changes seen during the programme. Wilcoxon signed rank and Friedman tests were used to examine whether there were significant differences in variables across time points.

In the longitudinal analysis, all variable scores at entry were force entered into a simple linear regression model and used to predict scores of the same variable at exit; standardised residual scores were retained. The same procedure was used when examining if exit predicted six month follow-up. Residual scores were used as indicators of change in each variable and were used in subsequent analyses. Change scores (residuals) were force entered into linear regression analyses to explore: (1) change in psychological needs predicting change in behavioural regulations (entry to exit) and (2) change in psychological needs and behavioural regulations (entry to exit) predicting change in psychological outcomes and habitual activity (entry to exit and exit to follow-up).

Results

Descriptive statistics

Table 1 shows how competence and relatedness satisfaction both significantly increased from entry to exit with levels of competence remaining significantly higher at six months than at entry. In contrast, relatedness satisfaction significantly decreased from exit to six month follow-up, resulting in significantly lower relatedness satisfaction at six months post-programme than at entry. No changes were observed in autonomy satisfaction. External regulation significantly decreased from entry to exit and remained significantly different from entry to six months; however, intrinsic motivation remained unchanged during the programme but then significantly decreased from baseline to six months.

All psychological outcomes improved significantly from entry to exit, with improvements in physical self-worth and the PCS of the SF-36v2 maintained at six months. Significant increases in habitual physical activity were seen at programme exit and were maintained at six month follow-up.

Table 1. Entry, exit and six month scores for psychological need satisfaction, behavioural regulation and psychological well-being.

	Score range	<i>N</i>	Entry mean \pm SD	Exit mean \pm SD	Six month mean \pm SD
Autonomy	0–12	184	8.24 \pm 2.66	8.43 \pm 2.61	8.23 \pm 2.70
Competence	0–12	184	8.90 \pm 2.40	9.58* \pm 2.25	9.09*** \pm 2.41
Relatedness	0–12	184	10.81 \pm 1.81	11.01* \pm 1.87	10.54*** \pm 2.07
Amotivation	0–4	200	.20 \pm .54	.18 \pm .47	.21 \pm .55
External regulation	0–4	200	.71 \pm .94	.58* \pm .86	.62* \pm .91
Introjected regulation	0–4	200	1.56 \pm 1.10	1.62 \pm 1.11	1.55 \pm 1.11
Identified regulation	0–4	200	3.06 \pm .75	3.08 \pm .70	3.06 \pm .75
Intrinsic motivation	0–4	200	3.02 \pm .85	3.09 \pm .82	2.97** \pm .91
Habitual physical activity	2–12	247	5.42 \pm 1.51	5.80* \pm 1.50	5.58*** \pm 1.56
Anxiety	0–21	255	6.91 \pm 4.65	6.45* \pm 4.39	6.80** \pm 4.60
Depression	0–21	254	4.93 \pm 3.67	4.61* \pm 3.53	4.78* \pm 3.71
Physical self-worth	6–24	226	13.58 \pm 3.94	14.23* \pm 3.77	13.83*** \pm 3.84
PCS of SF-36v2	0–100	258	38.67 \pm 10.13	40.22* \pm 10.34	39.11*** \pm 10.24
MCS of SF-36v2	0–100	258	44.74 \pm 12.50	46.39* \pm 12.41	44.94** \pm 12.79

*significant difference from entry $p < .05$.

**significant difference from exit $p < .05$.

Table 2. Correlation matrix of residual scores.

	Change scores (residuals) from entry-exit programme							
	Autonomy	Competence	Relatedness	Amotivation	External regulation	Introjected regulation	Identified regulation	Intrinsic motivation
Change scores (residuals) from entry – exit of programme								
Amotivation	-.235**	.150*	.017					
External	-.185*	-.139	.075					
Introjected	.087	.096	.168*					
Identified	.085	.142	.204**					
Intrinsic	.259**	.312**	.154*					
Anxiety	-.137	-.177*	-.091*	.132	.078	-.187*	-.078	-.008
Depression	-.017	-.224**	-.166	.067	-.021	-.111	-.128	-.092
Physical SW	-.025	.160	.104	-.009	-.246**	-.005	.156	.191*
PCS of SF-36v2	.151	.269**	.138	-.041	-.190*	.149	.110	.272**
MCS of SF-36v2	.208*	.199**	.218**	-.023	-.013	.134	.094	-.094
Habitual PA	-.001	-.264**	.085**	-.052	-.116	.159*	.113	.122
Anxiety	.161*	.095	-.004	-.104	.029	.178*	.046	.011
Depression	.039	.126	.048	-.053	.047	.053	.116	.120
Physical SW	-.032	-.032	-.047	-.144	-.050	.105	-.042	-.073
MCS of SF-36v2	-.261**	-.066	-.167	.027	.029	.135	.057	-.120
PCS of SF-36v2	-.067	-.138	-.088	-.035	-.045	-.128	-.110	-.154
Habitual PA	.073	-.023	.048	-.033	.017	.095	.256**	.256**

Note: Habitual PA – Habitual physical activity; Physical SW – Physical self-worth.
 * $p < .05$; ** $p < .01$.

Inferential statistics

Table 2 shows the correlation matrix between all residual scores used in subsequent regression models. Table 3 shows how changes in psychological needs predicted changes in outcome variables during the programme. Increases in autonomy satisfaction significantly predicted decreases in amotivation and external regulation and increases in intrinsic motivation. Increases in competence satisfaction predicted an increase in intrinsic motivation, physical well-being and habitual physical activity and a decrease in depression. Finally, increases in relatedness satisfaction predicted increases in identified regulation and mental well-being.

Table 4 shows how changes in behavioural regulation (entry to exit) predicted changes seen in psychological well-being (entry to exit). A decrease in external

Table 3. Changes in psychological need satisfaction from entry to exit predicting changes in behavioural regulation, well-being and habitual physical activity from entry to exit.

	<i>N</i>	Adj. <i>R</i> ²	<i>F</i>	<i>B</i>	SE	β
Amotivation	176	.061	4.777*			
Autonomy				-.212	.070	-.226**
Competence				-.131	.073	-.139
Relatedness				.099	.072	.106
External regulation	175	.049	4.019**			
Autonomy				-.172	.071	-.181*
Competence				-.143	.074	-.151
Relatedness				.141	.073	.149
Introjected regulation	175	.017	1.992			
Identified regulation	175	.034	3.041*			
Autonomy				.042	.074	.044
Competence				.081	.077	.083
Relatedness				.167	.076	.172*
Intrinsic motivation	175	.123	9.098**			
Autonomy				.175	.065	.195**
Competence				.237	.067	.263**
Relatedness				.041	.067	.045
Anxiety	155	.025	2.341			
Depression	155	.033	2.771*			
Autonomy				.044	.078	.046
Competence				-.223	.081	-.230**
Relatedness				.001	.077	.001
Physical self-worth	147	.027	2.365			
PCS of SF-36V2	129	.079	4.641**			
Autonomy				.109	.086	.108
Competence				.259	.086	.249**
Relatedness				.143	.111	.110
MCS of SF-36V2	129	.088	5.127**			
Autonomy				.192	.099	.166
Competence				.188	.099	.163
Relatedness				.286	.128	.190*
Habitual physical activity	154	.054	3.886*			
Autonomy				-.041	.070	-.047
Competence				.225	.069	.269**
Relatedness				.009	.068	.011

* $p < .05$; ** $p < .01$.

Table 4. Changes in behavioural regulation from entry to exit predicting changes in well-being and habitual physical activity from entry to exit.

	<i>N</i>	Adj. R^2	<i>F</i>	<i>B</i>	SE	β
Anxiety	174	.035	2.256			
Depression	173	-.003	.903			
Physical self-worth	159	.059	2.972*			
Amotivation				.101	.082	.102
External regulation				-.247	.089	-.230**
Introjected regulation				-.027	.085	-.027
Identified regulation				-.031	.102	-.031
Intrinsic motivation				.193	.105	.181
PCS of SF-36v2	133	.085	3.459**			
Amotivation				.140	.100	.133
External regulation				-.197	.110	-.157
Introjected regulation				.020	.088	.119
Identified regulation				-.092	.117	-.084
Intrinsic motivation				.398	.139	.319**
MCS of SF-36v2	133	.025	1.670			
Habitual physical activity	171	.034	2.180			

* $p < .05$; ** $p < .01$.

regulation predicted an increase in physical self-worth, whilst increases in intrinsic motivation predicted increases in SF-36v2-PCS.

The results of analyses exploring whether changes in psychological needs and behavioural regulation that were experienced during the programme predicted psychological well-being and habitual physical activity from exit to follow-up demonstrated that changes in autonomy satisfaction from entry to exit negatively predicted changes in the mental component of quality of life (Adj. $R^2 = .068$, $F(128) = 4.122$, $p < .01$; standardised $\beta = -.244$, $p < .01$), whilst changes in intrinsic motivation from entry to exit positively predicted changes in habitual physical activity from exit to six months (Adj. $R^2 = .075$, $F(170) = 3.774$, $p < .01$; standardised $\beta = -.219$, $p < .05$).

Discussion

This research employed SDT as a framework to explore how psychological need satisfaction and behavioural regulation relate to change in CR patients' psychological well-being and physical activity following an exercise-based CR programme. The hypothesis was supported. Specifically, increased autonomy satisfaction during the programme predicted decreased amotivation and external regulation and increased intrinsic motivation. It appears that autonomy was central to internalising behavioural regulations for exercise. However, satisfaction of this psychological need had no direct benefit for psychological well-being.

Both competence and relatedness satisfaction predicted behavioural regulation and psychological and behavioural outcomes in the expected direction. Competence satisfaction predicted an increase in intrinsic motivation, physical quality of life and habitual physical activity, as well as a decrease in depression during the supervised CR exercise programme. It appears that following the potential disruption to physical ability and daily functioning resulting from a cardiac event or disease (Hobbs et al., 2002), whether an individual subsequently feels competent to exercise contributes to their enjoyment of CR, their perception of their physical well-being and depression level. It also makes

intuitive sense that an individual's perceived ability to exercise is likely to positively influence their levels of activity (Rhodes & Nigg, 2011).

This lends support to findings by Russell and Bray (2009) who identified competence satisfaction as a key predictor of self-determined motivation in CR. Given that competence satisfaction is modifiable, for instance, by providing positive feedback (Ryan, Patrick, Deci, & Williams, 2008), this finding identifies a potential mechanism for reducing depression, considered to be a key contributor to secondary cardiac events (Frasure-Smith & Lespérance, 2005, 2008), as well as encouraging self-determined motivation to engage in CR and physical activity.

Increases in relatedness satisfaction significantly predicted increased identified regulation and mental well-being. Thus, feeling connected to the exercise environment may contribute to the internalisation process whereby individuals begin to identify with the values of exercise. This sense of support is key to enhancing psychological aspects of quality of life.

Becoming less externally regulated from entry to exit predicted increases in physical self-worth. This may stem from increased pride in physical achievements given the opportunity to regain ownership of one's behaviour, feeling less controlled by external sources and actively participating in risk factor modification. This relationship between external regulation and physical self-worth therefore suggests the importance of encouraging CR participants to be more self-determined during programme delivery.

Finally, increasing intrinsic motivation from entry to exit predicted improvements in the quality of life-PCS over this period, supporting the SDT proposal that self-determined motivation is necessary for increased well-being (Deci & Ryan, 2002). A possible explanation for this relationship is that participants who became more self-determined during the programme adhered more and benefitted from the exercise, and thus, they perceived an improvement in their physical well-being. Alternatively, individuals who enjoyed the exercise might have been more likely to acknowledge the benefits accrued.

There were no associations between changes in behavioural regulation from entry to exit and changes in well-being at six month follow-up. However, increased intrinsic motivation during the programme predicted subsequent increases in habitual physical activity at six month follow-up. As the aim of CR is to encourage long-term behaviour change, this finding indicates the importance of helping patients to feel self-determined and enjoy their exercise sessions in achieving this aim. Increased autonomy satisfaction significantly predicted decreases in the mental aspects of quality of life from exit to six months. This is an unexpected finding and suggests that those who became more autonomous during the programme were less likely to adapt well psychologically in the sixth months following the structured programme. It may be that those individuals with higher levels of autonomy satisfaction and who were therefore participating through volitional choice felt despondent when the formal element of the programme ended thus having a negative impact on their mental well-being.

The lack of other significant longitudinal relationships between changes in need satisfaction and psychological outcomes in the six month post-programme suggests that effects might be context specific. Thus, changes experienced could well have little impact after formal delivery of the programme where structured support of need satisfaction is available for participants. In this case, social and environmental factors such as access to facilities and social support from family and friends might better explain subsequent changes in psychological need satisfaction.

However, limitations of the current study could also contribute to these non-significant effects. Some variables were not normally distributed which should be considered when interpreting regression analyses. Although attributable to the clinical sample including a higher proportion of participants with low well-being scores compared with those typically seen in a non-clinical sample (Vickers, 2007), this is likely to have resulted in reduced power.

Despite this, this study has advanced the current literature by demonstrating how changes in components of SDT might help to explain changes in psychological well-being during a CR programme and in physical activity following the programme. Competence satisfaction and self-determined regulation appear to contribute significantly to improved well-being and activity within this context, lending support to previous research in CR. Although the study demonstrated that the CR programme was effective in improving individuals' psychological need satisfaction, the programme was not designed with this explicit goal in mind. Therefore, future research would benefit from examining the effects of an SDT-based exercise intervention on psychological and behavioural outcomes in similar clinical populations.

Funding

This work was supported by Big Lottery Fund.

ORCID

Rachel Jane Rahman  <http://orcid.org/0000-0003-2191-8059>

References

- Baecke, J. A. H., Burema, J., & Frijters, J. E. R. (1982). A short questionnaire for the measurement of habitual physical activity in epidemiological studies. *American Journal of Clinical Nutrition*, *36*, 936–942.
- Balady, G., Fletcher, B., & Froelicher, E. (1994). Cardiac rehabilitation programs: A statement for health care professionals from the American Heart Association. *Circulation*, *90*, 1602–1610.
- Bjarnason-Wehrens, B., Bott, D., Benesch, L., Bischoff, K. O., Buran-Kilian, B., Gysan, D., ... Sauer, G. (2007). Long-term results of a three-week intensive cardiac out-patient rehabilitation program in motivated patients with low social status. *Clinical Research in Cardiology*, *96*, 77–85. doi:10.1007/s00392-007-04610-0
- British Heart Foundation. (2010). *The national audit of cardiac rehabilitation; annual statistics report 2010*. Northampton: Author.
- Deci, E. L., & Ryan, R. M. (1985). *Intrinsic motivation and self-determination in human behavior*. New York, NY: Plenum.
- Deci, E. L., & Ryan, R. M. (2002). An overview of self-determination theory: An organismic-dialectical perspective. In E. L. Deci & R. M. Ryan (Eds.), *Handbook of self-determination research* (pp. 3–33). Rochester, NY: University of Rochester Press.
- Florindo, A. A., do Rosario Dias, M., & Latorre, O. (2003). Validation and reliability of the Baecke questionnaire for the evaluation of habitual physical activity in adult men. *Revista Brasileira de Medicina do Esporte*, *9*, 1517–8692. doi:10.1590/s1517-86922003000300002
- Fox, K. R., & Corbin, C. B. (1989). The physical self-perception profile: Development and preliminary validation. *Journal of Sport and Exercise Psychology*, *11*, 408–430.
- Frasure-Smith, N., & Lespérance, F. (2005). Reflections on depression as a cardiac risk factor. *Psychosomatic Medicine*, *67*, S19–S25. doi:10.1097/01.psy.0000162253.07959.db
- Frasure-Smith, N., & Lespérance, F. (2008). Depression and anxiety as predictors of 2-year cardiac events in patients with stable coronary artery disease. *Archives of General Psychiatry*, *65*, 62–71. doi:10.1001/archgenpsychiatry.2007.4

- Hobbs, F. D. R., Kenkre, J. E., Roalfe, A. K., Davis, R. C., Hare, R., & Davies, M. K. (2002). Impact of heart failure and left ventricular systolic dysfunction on quality of life. A cross-sectional study comparing common chronic cardiac and medical disorders and a representative adult population. *European Heart Journal*, *23*, 1867–1876. doi:10.1053/euhj.2002.3255
- Jenkinson, C., Stewart-Brown, S., Petersen, S., & Paice, C. (1999). Assessment of the SF-36 version 2 in the United Kingdom. *Journal of Epidemiology and Community Health*, *53*, 46–50. doi:10.1136/jech.53.1.46
- Markland, D., & Tobin, V. (2004a). A modification to the behavioural regulation in exercise questionnaire to include an assessment of amotivation. *Journal of Sport and Exercise Psychology*, *26*, 191–196.
- Markland, D., & Tobin, V. (2004b). On the relationships among perceived environmental supportiveness, psychological need satisfaction and intrinsic motivation: A comparison of alternative models. International Self-Determination Theory Conference II, Ottawa, Canada.
- Moorey, S., Greer, S., Watson, M., Gorman, C., Rowden, L., Tunmore, R., ... Bliss, J. (1991). The factor structure and factor stability of the hospital anxiety and depression scale in patients with cancer. *The British Journal of Psychiatry*, *158*, 255–259. doi:10.1192/bjp.158.2.255
- National Institute for Clinical Excellence. (2013). *MI: Secondary prevention in primary and secondary care for patients following a myocardial infarction*. London: NICE Clinical guidance 172.
- Ng, J. Y. Y., Ntoumanis, N., Thogersen-Ntoumani, C., Deci, E. L., Ryan, R. M., Duda, J. L., & Williams, G. C. (2012). Self-determination theory applied to health contexts: A meta-analysis. *Perspectives on Psychological Science*, *7*, 325–340.
- Rahman, R. J., Thogersen-Ntoumani, C., Thatcher, J., & Doust, J. (2011). Changes in need satisfaction and motivation orientation as predictors of psychological and behavioural outcomes in exercise referral. *Psychology & Health*, *26*, 1521–1539. doi:10.1080/08870446.2010.538849
- Rhodes, R. E., & Nigg, C. R. (2011). Advancing physical activity theory. *Exercise and Sport Sciences Reviews*, *39*, 113–119.
- Rouse, P. C., Ntoumanis, N., Duda, J. L., Jolly, K., & Williams, G. C. (2011). In the beginning: Role of autonomy support on the motivation, mental health and intentions of participants entering an exercise referral scheme. *Psychology & Health*, *26*, 729–749. doi:10.1080/08870446.2010.492454
- Russell, K. L., & Bray, S. R. (2009). Self-determined motivation predicts independent, home based, exercise following cardiac rehabilitation. *Rehabilitation Psychology*, *54*, 150–156. doi:10.1037/a0015595
- Ryan, R. M., Patrick, H., Deci, E. L., & Williams, G. C. (2008). Facilitating health behaviour change and its maintenance. Interventions based on self-determination theory. *The European Health Psychologist*, *10*, 2–5.
- Sweet, S. N., Tulloch, H., Fortier, M. S., Pipe, A. L., & Reid, R. D. (2011). Patterns of motivation and ongoing exercise activity in cardiac rehabilitation settings: A 24-month exploration from the teach study. *Annals of Behavioral Medicine*, *42*, 55–63.
- Vickers, A. J. (2007). If the normal distribution is so normal, how come my data never are? *Medscape Business of Medicine*. Retrieved May 15, 2007 from <http://www.medscape.com/viewarticle/556012>
- Wade, J. E., Snow, K. K., Kosinski, M., & Gandek, B. (1993). *SF-36® health survey: Manual & interpretation guide*. Lincoln, RI: QualityMetric Incorporated.
- Wade, J. E., Snow, K. K., Kosinski, M., & Gandek, B. (2002). *SF-36® health survey: Manual & interpretation guide*. Lincoln, RI: QualityMetric Incorporated.
- Wilson, P. M., Rodgers, W. M., Blanchard, C. M., & Gessell, J. G. (2003). The relationship between psychological needs, self-determined motivation, exercise attitudes, and physical fitness. *Journal of Applied Social Psychology*, *33*, 2373–2392. doi:10.1111/j.1559-1816.2003.tb01890.x
- Yohannes, A. M., Doherty, P., Bundy, C., & Yalfani, A. (2010). The long term benefits of cardiac rehabilitation on depression, anxiety, physical activity and quality of life. *Journal of Clinical Nursing*, *19*, 2806–2813. doi:10.1111/j.1365-2702.2010.03313.x
- Zigmond, A. S., & Snaith, R. P. (1983). The hospital anxiety and depression scale. *Acta Psychiatrica Scandinavica*, *67*, 361–370. doi:10.1111/j.1600-0447.1983.tb09716.x